

A CHECK-LIST OF THE NEUROPTERA- PLANIPENNIA OF THE USSR FAR EAST, WITH SOME TAXONOMIC REMARKS

V. N. MAKARKIN

Institute of Biology and Pedology, Far Eastern Division of the USSR Academy of Sciences, Vladivostok-22, USSR

(Received 15 May, 1989)

Seventy-seven species of the Far Eastern Neuroptera are listed. *Chrysopa cognata* is considered to be a separate species from *Ch. septempunctata*. *Wesmaelius pseudofurcatus*, *Mantispa kononenkoi*, and *Euroleon sinicus* are synonymized with *W. furcatus*, *M. styriaca*, and *E. coreanus* respectively. *Drepanepteryx pleshanovi* is now considered as a colour form of the *D. punctata*. With 5 figures.

Up to now 67 species of Neuroptera-Planipennia have been recorded from the Far East of the USSR. In this paper I add further 11 species, from which three ones are new to the USSR fauna (*Wesmaelius asiaticus*, *Dendroleon jezoensis* and *Glenuroides japonicus*); *Drepanepteryx pleshanovi* MAKARKIN, 1985 is now considered to be a colour form of *D. punctata* OKAMOTO. Thus 77 species are included in this checklist. Of these 2 species (*Micromus multipunctatus* and *Chrysopa sapporensis*) are unknown to me in nature. The cited literature only include the works contained the records of the USSR Far Eastern Neuroptera. New records are asterisked (*). The USSR Far East regions are shown in Figure 1.

A b b r e v i a t i o n s — Am = Amur province, Ch = Chukotka, Kh = Khabarovsk province, Km = Kamchatka, Kr = Kurile islands, Mg = Magadan province, N Kh = North of Khabarovsk province, Pr = Primorje (Maritime province), Sa = Saghalien.

OSMYLIDAE

1. **Osmylus (O.) hyalinatus** MACLACHLAN, 1875 — KUWAYAMA 1936: 109, 1956: 79, 1967: 65 (*Plethosmylus*); MAKARKIN 1985c: 39, figs 8, 17, 25. — Distribution: Kr (Kunashir Is.), Sa. Japan.
2. **Osmylus (O.) decoratus** NAKAHARA, 1913 — MAKARKIN 1985c: 41, figs 2, 12, 13, 26. *hyalinatus* (nec MACLACHLAN): NAVÁS 1912: 421. — Distribution: Kh, Kr (Kunashir Is.), Pr. Japan.
3. **Osmylus (O.) pryeri** MACLACHLAN, 1875 — MAKARKIN 1985c: 41, figs 11, 15, 23. — Distribution: Kr (Kunashir Is.) Japan.
4. **Osmylus (Plesiosmylus) tessellatus** MACLACHLAN, 1875 — Kuwayama 1936: 108, 1956: 79, 1967: 65; MAKARKIN 1985c: 41, figs 7, 16, 28. — Distribution. Kr (Kunashir Is.) Japan.
5. **Lysmus harmandinus** (NAVÁS, 1910) — KUWAYAMA 1924: 115, 1936: 108 (*Eososmylus*), 1956: 79, 1967: 65; MAKARKIN, 1985c: 45, figs 3, 6, 21, 29. *kuriensis* KUWAYAMA, 1956: 20, fig. 2, 1967: 65. *flavicornis* (nec MACLACHLAN): MATSUMURA 1911: 15 (*Osmylus*); KUWAYAMA 1936: 104 (*Heliosmylus*).

SISYRIDAE

6. *Sisyra terminalis* CURTIS, 1834 — DOROKHOVA 1987: 48; ZAKHARENKO 1988: 766, fig. 23. — Distribution: Kh, Pr. Europe.

7. *Sisyra nikkoana* (NAVÁS, 1910) — KUWAYAMA 1967: 65; ZAKHARENKO 1988: 767, figs 17–21. — Distribution: *Kh (55 km east of Birobidzhan), Kr (*Kunashir, Is. Iturup Is.), Pr, *Sa. Japan.

DILARIDAE

8. *Dilar septentrionalis* NAVÁS, 1912 — NAVÁS 1912: 420; ZAKHARENKO 1988: 764, figs 1–7. — Distribution: Pr. — China.

MANTISPIDAE

9. *Mantispa styriaca* (PODA, 1761) — ZAKHARENKO, 1987a: 622. — *kononenkoi* MAKARKIN, 1985b: 620 figs a–d, syn. n. — Distribution: Pr. Europe, Asia Minor, Armenia, North Iran, Middle Asia, Mongolia, China.




Fig. 1. Far Eastern regions: Am = Amur province, Ch = Chukotka, Kh = Khabarovsk province, Km = Kamchatka, Kr = Kurile Islands, Mg = Magadan province, N Kh = North of Khabarovsk province, Pr = Prymorje, Sa = Saghalien

10. **Mantispa japonica** MACLACHLAN, 1875 — MAKARKIN 1985b: 622, fig. e—f. — Distribution: Pr. Japan, China, Korea. — Remarks: It is possible that this species is a synonym of *M. perla* PALLAS or *M. lobata* NAVÁS.

11. **Eumantispa harmandi** (NAVÁS, 1909) — MAKARKIN 1985b: 622; ZAKHARENKO 1987a: 624, figs 2, 13. — Distribution: Pr. Japan, Korean Peninsula.

CHrysopidae

12. **Chrysopa (Ch.) perla** (LINNAEUS, 1758) — NAVÁS 1912: 419; PLESHANOV 1974: 189; DOROKHOVA 1980: 21; MAKARKIN 1985f: 57, fig. 4. — Distribution: Kh, *N Kh, Sa. Palaearctic Region.

13. **Chrysopa (Ch.) intima** MACLACHLAN, 1893 — KUWAYAMA 1924: 110, 1936: 108, 1956: 81, 1967: 65; PLESHANOV 1974: 189; DOROKHOVA 1979: 107; MAKARKIN 1985e: 91, 1985f: 57, fig. 5. var. *fracta* NAVÁS, 1912: 419 (*Chrysopa perla*). (?) *perla* (nec LINNAEUS): MATSUMURA, 1911: 14. — Distribution: Am, Kh, Km, Kr (Iturup Is., Kunashir Is.), Pr, Sa. *Jakutia, Siberia, Mongolia, China, Japan, Korea.

14. **Chrysopa (Ch.) formosa** BRAUER, 1850 — PLESHANOV 1974: DOROKHOVA 1979: 107; SHUVAKHINA 1980a: 22; MAKARKIN 1985e: 91, 1985f: 57, fig. 6. — Distribution: Am, Kh, Km, *Kr (Kunashir Is.), Pr, *Sa. Palaearctic Region.

15. **Chrysopa (Ch.) abbreviata** CURTIS, 1854 — Distribution: *N Kh (District of Okhotsk). Europe, Siberia, Middle Asia, Mongolia.

16. **Chrysopa (Ch.) commata** KIS et ÚJHELYI, 1965 — MAKARKIN 1985e: 91, 1985f: 59, figs 1, 9. — Distribution: Am, Kh, Kr (Kunashir Is.), Pr. Europe, Armenia, North Iran, Siberia.

17. **Chrysopa (Ch.) sapporensis** OKAMOTO, 1914 — KUWAYAMA 1956: 81. — *phyllochroma* (nec WESMAEL): KUWAYAMA 1967: 66. — Distribution: Kr (Iturup Is.). Japan. — Remark: *Ch. sapporensis* was considered to be a synonym of *Ch. phyllochroma*. However, TSUKAGUCHI (1985) found that the latter species do not occur in Japan. It is possible that *Ch. sapporensis* and *Ch. commata* refer to the same species.

18. **Chrysopa (Ch.) perplexa** MACLACHLAN, 1887 — DOROKHOVA 1979: 108; MAKARKIN 1985e: 91, 1985f: 60, figs 3, 8. — Distribution: Am, Kh, Pr. East Siberia, China, Japan, Korea.

19. **Chrysopa (Ch.) phyllochroma** WESMAEL, 1841 — (?) PLESHANOV 1974: 188; MAKARKIN 1985f: 61, figs 2, 10. — Distribution: Am, ?Pr. Palaearctic Region.

20. **Chrysopa (Parachrysopa) septempunctata** WESMAEL, 1841 — KUWAYAMA 1924: 111, fig. 3; PLESHANOV 1974: 188 (partim); DOROKHOVA 1979: 111 (partim); SHUVAKHINA 1980b: 23 (partim); MAKARKIN 1985: 58, fig. 7 (partim). — Distribution: Am, Kh, Km, Pr, Sa. Palaearctic Region.

21. **Chrysopa (Parachrysopa) cognata** MACLACHLAN, 1867 — NAVÁS 1912: 419. *septempunctata* (nec WESMAEL): PLESHANOV 1974: 188 (partim); DOROKHOVA 1979: 111 (partim); SHUVAKHINA 1980b: 23 (partim); MAKARKIN 1985e: 91; 1985f: 58, fig. 7 (partim). — Distribution: *Am, *Kh, Pr. China, Korea, Japan, Cambodia. — Remark. This species was described from Cambodia, China and Japan, but for a long time it was considered as a synonym of *Ch. septempunctata*. However, *Ch. cognata* differs from the latter species externally as well as in male genitalia structures as follows:


1 (2) Body wholly green; head mainly with 7 spots: on genae (2), on clypeus (2), under antennae (2), and between antennae (1). Entoprocessus long, extending out of hind margin of gonarcus (Fig. 2) **Ch. septempunctata**

2 (1) Body green dorsally with median yellow stripe; head mainly with 4—5 spots: on clypeus (2), under antennae (2), and between antennae (1 or absent). Entoprocessus short and curved (Fig. 3) **Ch. cognata**

22. **Chrysopa (Cunctochrysa) albolineata** KILLINGTON, 1935 — MAKARKIN 1985e: 91, 1985f: 61, fig. 11. — Distribution: Pr. Europe, Anatolia, Armenia, Iran, Afghanistan, Japan, Korea.

23. **Chrysopa (?Aperthochrysa) joannisi** NAVÁS, 1910 — MAKARKIN, 1985d: 48, figs 1—5; 1985f: 61, fig. 12. — Distribution: Pr. China.

24. **Chrysopa (?Chrysoperla) cognatella** OKAMOTO, 1914 — KUWAYAMA 1962: 366; MAKARKIN 1985d: 48, figs 6—8, 1985f: 61, fig. 13. — Distribution: Pr, Sa. Japan, Korea, China (including Taiwan).



Figs 2—3. Male gonarcus and pseudopenis of *Chrysopa septempunctata* WESM. (2) and *Ch. cognata* McL. (3), lateral view

25. *Chrysoperla carnea* (STEPHENS, 1836) — PLESHANOV 1974: 187; MAKARKIN 1985f: 61, fig. 14. — Distribution: Am, Kh, Pr. Holarctic Region.

26. *Chrysoperla nippensis* (OKAMOTO, 1914) — MAKARKIN 1985e: 91, 1985f: 61, fig. 15. *sinica* (? nec TJEDER, 1936): DOROKHOVA 1979: 110 (*Chrysopa*). — Distribution: Kh, Kr (Kunashir Is.), Pr. China, Japan, Korea.

27. *Chrysotropia ciliata* (WESMAEL, 1841) — PLESHANOV 1974: 187; DOROKHOVA 1979: 106; MAKARKIN 1985e: 91, 1985f: 62, fig. 19. — Distribution: Am, Kh, Kr (Kunashir Is.), Pr. Sa. Palaearctic Region.

28. *Mallada prasina* (BURMEISTER, 1839) — KUWAYAMA 1967: 66 (*Chrysopa*); PLESHANOV 1974: 187 (*Chrysopa*); DOROKHOVA 1979: 109, 1981: 5 (*Chrysopa*); MAKARKIN 1985e: 91, 1985f: 61, fig. 16 (*Anisochrysa*). *sachalinensis* MATSUMURA, 1911: 14 (*Chrysopa*); KUWAYAMA 1924: 111, 1956: 82 (*Chrysopa*). — Distribution: Am, Kh, Kr (Kunashir Is.), Shikotan Is., Pr, Sa. Palaearctic Region.

29. *Mallada laurae* (MAKARKIN, 1985) comb. n. — MAKARKIN 1985d: 49, figs 9—14, 1985f: 62, fig. 17 (*Anisochrysa*). — Distribution: Kh, Kr (Kunashir Is.), Pr, *Sa.

30. *Mallada ussuriensis* (MAKARKIN, 1985) comb. n. — MAKARKIN 1985d: 51, figs 15—20 1985e: 91, 1985f: 62, fig. 18 (*Anisochrysa*). — Distribution: Kh, Pr.

31. *Nineta vittata* (WESMAEL, 1841) — NAVÁS 1925: 1; KUWAYAMA 1924: 109, 1936: 108, 1956: 81 (*Chrysopa*), 1967: 66 (*Chrysopa*); PLESHANOV 1974: 187; DOROKHOVA 1979: 108; MAKARKIN 1985f: 62, fig. 20. *inornata* MATSUMURA 1911: 14 (*Chrysopa*). — Distribution: Kh, Km, Kr (Iturup Is., Kunashir Is., Shikotan Is.), Pr, Sa. Palaearctic Region.

32. *Nineta carinthiaca* (HÖLZEL, 1965) — MAKARKIN 1985d: 52, figs 21—24, 1985e: 91, 1985f: 62, fig. 21. — Distribution: Am, Kh, Pr. Europe, Siberia.


HEMEROBIIDAE

33. *Hemerobius (H.) humulinus* LINNAEUS, 1758 — NAVÁS 1912: 419, 1925: 2; ESBEN-PETERSEN 1921: 40; KUWAYAMA 1936: 109, 1962: 353; MAKARKIN, 1985a: 163, figs 16, 32; 1985e: 91. ? *shikotanus* KUWAYAMA, 1956: 77, fig. 1; 1967: 65. — Distribution: Am, Kh, Km, Kr (Kunashir Is., ? Shikotan Is.), Mg, Pr, Sa. Holarctic Region. — Remarks: An examination of the female holotype of *H. shikotanus* will decide whether it is identical with *H. humulinus* or *H. japonicus*?

34. *Hemerobius (H.) japonicus* NAKAHARA, 1915 — KUWAYAMA 1956: 77, 1967: 65; MAKARKIN 1985a: 164, figs 17, 36; 1985e: 91. — Distribution: Kr (Kunashir Is.), Pr. Japan.

35. *Hemerobius (H.) simulans* WALKER, 1853 — MAKARKIN 1985a: 165, figs 18, 44, 1985e: 91. — *piceus* NAVÁS 1925: 2, fig. 1. — Distribution: Am, Kh, N Kh, Km, Mg, Pr, Sa. Holarctic Region.

36. *Hemerobius (H.) fujimotoi* NAKAHARA, 1960 — Distribution: *Am, *Kh, *Sa. Caucasus, Siberia, Japan.
37. *Hemerobius (H.) fenestratus* TJEADER, 1932. — MAKARKIN, 1985a: 166, figs 12, 33. — Distribution: *Am, *Kh, *Km, Kr (Kunashir Is.), Pr, Sa. Europe, Mongolia, Siberia, ? Japan.
38. *Hemerobius (H.) atrifrons* MACLACHLAN, 1868 — MAKARKIN 1985a: 166, figs 15, 29. — Distribution: Am, Kh, *N Kh, Km, Mg, Sa. Europe, Mongolia, China, Japan.
39. *Hemerobius (H.) nitidulus* FABRICIUS, 1777 — NAVÁS 1925: 2; MAKARKIN, 1985a: 165, figs 11, 31. — Distribution: Km, Kr (Paramushir Is.), Mg. Europe, Caucasus, Siberia, Mongolia.
40. *Hemerobius (H.) stigma* STEPHENS, 1836 — MAKARKIN, 1985a: 166, figs 15, 29. — Distribution: Am, *N Kh, Km, Mg, *Pr, Sa. Holarctic Region.
41. *Hemerobius (Brauerobius) marginatus* Stephens, 1836 — ESBEN-PETERSEN 1921: 40; NAVÁS 1925: 2; MAKARKIN 1985a: 167, figs 27, 39. — Distribution: Kh, *N Kh, Km, Mg, Pr, Sa. Palaearctic Region. — Remarks: In the northern part of its range, this species is considerably smaller and darker. Such specimens from Finland were named by MEINANDER (1962) as *H. m. lapponicus*. It occurs also in the Komi ASSR (ZAKHARENKO & SÉDYKH 1981), in Jakutia and in the northern part of the Far East and are possibly identical with *H. costalis* CARPENTER distributed in North America (including Alaska).
42. *Hemerobius (Brauerobius) tristriatus* KUWAYAMA, 1954 — KUWAYAMA 1954: 97; MAKARKIN 1985a: 167, figs 26, 37. — Distribution: *Am, Kh, Kr (Kunashir Is., Shikotan Is.), Pr, Sa. Japan.
43. *Hemerobius (Allemerobius) exoterus* NAVÁS, 1936 — MAKARKIN 1985a: 168, figs 24, 45; 1985e: 91. — Distribution: Pr. China. — Remarks: The subgenus *Monorobius* MAKARKIN, 1985 (type-species *H. lutescens*, by original designation) is a junior synonym of *Allemerobius* BANKS, 1940 (type-species *A. flaveolus* BANKS, 1940 by original designation) syn. n.). BANKS (1905) designated *H. lutescens* F. as genotype of *Mucropalpus* RAMBUR, 1842. However, *H. lutescens* in sensu RAMBUR, 1842 is identical with *H. humulinus* L., the type species of *Hemerobius* LINNAEUS, 1758 (HAGEN 1866, LERAUT 1980). Therefore, *Mucropalpus* is an objective synonym of *Hemerobius*.
44. *Hemerobius (Allemerobius) poppii* ESBEN-PETERSEN, 1921 — MAKARKIN 1985a: 168, figs 25, 43. — Distribution: *Am, *Km, *N Kh, Mg, *Sa. Siberia, China, Mongolia.
45. *Hemerobius (Allemerobius) subfalcatus* NAKAHARA, 1960 — MAKARKIN 1985a: 168, figs 20, 42. — Distribution: *Kr (Kunashir Is.), *Pr, Sa. Japan.



Figs 4—5. *Wesmaelius asiaticus* YANG, basal parts of the antennae of male (4) and female (5), lateral view

46. *Wesmaelius (W.) quadrifasciatus* (REUTER, 1894) — NAKAHARA 1956: 182; MAKARKIN 1986: 607, figs 13, 15. — Distribution: Am, Kh, *Km, *N Kh, Kr (Kunashir Is.), Mg, Sa. Palaearctic Region.

47. *Wesmaelius (W.) asiaticus* YANG, 1980 — Distribution: *Kh, *Pr. China. — Remarks: This species was described on the basis of a single female from North China. It is closely related to *W. quadrifasciatus*, but differs from it by the genitalia structures of both males and females as well as by an unusual shape of the male antennae (Figs 4—5).

48. *Wesmaelius (Kimmminsia) nervosus* (FABRICIUS, 1793) — HAGEN 1858: 130 (*Hemero-bius*); NAVÁS 1925: 3 (*Hemero-bius*). *betulinus* STRÖM, 1788; MAKARKIN 1986: 607, figs 3, 17. — Distribution: *Ch, Km, Kr (Kunashir Is.), N Kh, Pr, Sa. Holarctic Region.

49. *Wesmaelius (Kimmminsia) furcatus* (BANKS, 1935): = *pseudofurcatus* MAKARKIN, 1986: 608, figs 7, 9, 10, 25 (syn. n.). — Distribution: Mg, Altaj, Buryatia, North America. — Remarks: *W. pseudofurcatus* was based mainly on the subgenitale plate characters, but it is recently showed that this structure is very variable (KLIMASZEWSKI & KEVAN 1987).

50. *Wesmaelius (?Kimmminsia) lateralis* (NAVÁS, 1912) — NAVÁS 1912: 419, fig. 4 (*Boriomyia*); MAKARKIN 1986: 611, figs 1, 11, 12, 27. — Distribution: Kh, Mg, Sa. Altaj, Buryatia, Japan.

51. *Neuronema laminata* TJEDER, 1936 — *tjederi* KIMMINS, 1943; MAKARKIN 1986: 613, fig. 34, 44. *albostigma* (nec MATSUMURA): PLESHANOV, 1974: 186. *deltoides* (nec NAVÁS, 1910): NAVÁS 1912: 420 (*Ninga*); (?) KUWAYAMA 1924: 107 (*Ninga*). — Distribution: *Am, *Kh, Pr, Sa. China.

52. *Neuronema albostigma* (MATSUMURA, 1907) — MAKARKIN, 1986: 613, figs 35, 45. — Distribution: Kr (Kunashir Is.). Japan.

53. *Drepanopteryx phalaenoides* (LINNAEUS, 1758) — NAVÁS 1929: 36; KUWAYAMA 1962: 358; MAKARKIN 1985b: 91; 1986: 615, fig. 36. — Distribution: *Kh, *Km, *Kr (Shikotan Is.), Pr, Sa. Palaearctic Region.

54. *Drepanopteryx algida* (ERICHSON, 1851) — MAKARKIN, 1986: 616, fig. 38. — Distribution: Am, *N Kh, *Sa. Europe, Siberia, Mongolia.

55. *Drepanopteryx punctata* (OKAMOTO, 1905) — KUWAYAMA 1962: 357 (*Oedobius*); MAKARKIN 1986: 616, fig. zh. — Distribution: Kr (Kunashir Is.), Pr, Sa. Japan. — Remarks: *D. pleshanovi* differs from *D. punctata* in having a longitudinal brown stripe on the fore wings. However, it does not differ from the latter species by the genitalia structures of both males and females. I believe that *D. pleshanovi* is a colour form of *D. punctata*. This form occurs in the Primorje and Saghaliens.

56. *Micromus angulatus* (STEPHENS, 1836) — NAVÁS 1912: 420; KUWAYAMA 1962: 350 (*Eumicromus*); MAKARKIN 1985a: 169, fig. 48, 1985e: 91. *aphidivorus*: NAVÁS 1925: 3. — Distribution: Am, Kh, Km, Mg, Pr, Sa. Holarctic Region.

57. *Micromus paganus* (LINNAEUS, 1767) — NAVÁS 1925: 3; KUWAYAMA 1956: 78 (*Eumicromus*); MAKARKIN 1985a: 169, fig. 46. *alpinus* NAKAHARA, 1915; KUWAYAMA 1956: 78 (*Eumicromus*) — Distribution: Am, Kh, Km, Kr (Kunashir Is., Paramushir Is., Urup Is.), Sa, Pr. Palaearctic Region.

58. *Micromus variegatus* (FABRICIUS, 1793) — MAKARKIN 1985a: 168, fig. 47, 1985e: 91. — Distribution: Pr. Asia Minor, Caucasus, North Iran, South Kazakhstan, Japan.

59. *Micromus multipunctatus* MATSUMURA, 1907 — KUWAYAMA 1962: 348. *novitius* NAVÁS, 1910; KUWAYAMA 1924: 108. — Distribution: ?Sa. Japan, China (including Taiwan Is.). — Remarks: I have not seen any specimens of this species from the USSR. As it may be confused with *Paramicromus dissimilis*, its distribution in our country needs a confirmation.

60. *Paramicromus dissimilis* (NAKAHARA, 1915) — KUWAYAMA 1962: 351; MAKARKIN 1985a: 170, fig. 49. — Distribution: Kr (Iturup Is.), Kunashir Is.), Sa. Japan.

61. *Psectra diptera* (BURMEISTER, 1839) — MAKARKIN 1985e: 91, 1986: 613. — Distribution: Km, Kr (Kunashir Is.), Pr. Holarctic Region.

62. *Sympherobius manchuricus* NAKAHARA, 1960 — MAKARKIN 1985e: 91, 1986: 611, fig. 39. — Distribution: *Am, Pr. China.

63. *Sympherobius fuscescens* (WALLENGREN, 1863) — Distribution: *Kr (Kunashir Is.). Europe, Anatolia, Kazakhstan, Siberia, Japan (Hokkaido). — Remark: Two females were found in the Kunashir Island. They are darker than the European and Siberian specimens. The Hokkaido specimen (a male) is also darker (NAKAHARA 1971).

CONIOPTERYGIDAE

64. *Coniopteryx (C.) pygmaea* ENDERLEIN, 1906 — MEINANDER 1981: 107. — Distribution: Kr (Kunashir Is.), Shikotan Is., *Pr. Europe.
65. *Coniopteryx (C.) parthenia* (NAVÁS et MARCET, 1910) — Distribution: *Kh. Europe, Morocco, Asia Minor, Siberia, Mongolia.
66. *Coniopteryx (C.) aspoecki* KIS, 1967 — Distribution: *Pr. Roumania, Austria, Jakutia.
67. *Coniopteryx (C.) helvola* ZAKHARENKO, 1987 — ZAKHARENKO 1987: 76, figs 1—5. — Distribution: Pr.
68. *Semidalis aleyrodiformis* (STEPHENS, 1836) — Distribution: *Pr. Palaearctic Region, India, Thailand, Malaya, Taiwan Is.

ASCALAPHIDAE

69. *Libelloides sibiricus* (EVERSMANN, 1850) — KOZHANCHIKOV 1953: 432 (*Ascalaphus*); PLESHANOV 1974: 191 (*Ascalaphus*). Distribution: *Am, *Kh, Pr. South Siberian, China, Korea.

MYRMELEONTIDAE

70. *Acanthaclisis japonica* MACLACHLAN, 1875 — MAKARKIN 1984: 38. — Distribution: Pr. China, Japan, Korea.
71. *Myrmeleon (M.) formicarius* LINNAEUS, 1767 — KUWAYAMA 1962: 388; PLESHANOV 1974: 190; MAKARKIN 1984: 39. — Distribution: Kh, Pr., Sa. Palaearctic Region.
72. *Myrmeleon (Morter) bore* (TJEDER, 1941) — KUWAYAMA 1962: 388, 1967: 66 (*Grocus*); PLESHANOV 1974: 191 (*Grocus*); MAKARKIN 1984: 39. *formicarius* (nec LINNAEUS): KUWAYAMA, 1936: 107, 1956: 82. — Distribution: *N Kh (district of Okhotsk), Kr (Kunashir Is.), Pr, Sa. Palaearctic Region.
73. *Deutoleon lineatus* (FABRICIUS, 1798) — NAVÁS 1912: 418 (*Formicaleo*); PLESHANOV 1974: 140 (*Formicaleo*); MAKARKIN 1984: 39. — Distribution: Kh, Pr. East Europe, Kazakhstan, South Siberia, Mongolia, China, Korea.
74. *Euroleon polystipus* (GERSTAECKER, 1885) — GERSTAECKER 1885: 24 (*Myrmeleon*); MAKARKIN 1984: 39 (*Formicaleo*). Distribution: *Kh (Angun river), Pr, Sa. Siberia, Mongolia.
75. *Euroleon coreanus* OKAMOTO, 1926 — OKAMOTO 1926: 19, fig. 1. *sinicus* NAVÁS, 1930; HÖLZEL 1970a: 254, fig. 13—15, syn. n. *nostras* (nec FOURCROY): PLESHANOV 1974: 190. — Distribution: *Pr. Burjatia, Mongolia, China, Korea. — Remark: I have examined two specimens from the Primorje (Novokachalinsk and Novonezhino) and a number of specimens from the Burjatia. The latter specimens were determined by H. HÖLZEL as *E. sinicus*. On the other hand, they agree well with the description of *E. coreanus*. I think that *E. sinicus* is a synonym of *E. coreanus*, although H. HÖLZEL (1970b) proposed the synonymy of *E. coreanus* with *E. polystipus*. However, the latter species differs in many respects from both *E. coreanus* and *E. sinicus*.
76. *Dendroleon jezoensis* OKAMOTO, 1910 — Distribution: *Pr. Japan, China, Korea. — Remark: I have examined only one male collected by me at Borisovka river 27 July 1987.
77. *Glenuroides japonicus* (MACLACHLAN, 1867) — Distribution: *Pr. Japan, China (including Taiwan Is.), Korea. — Remark: I have studied three specimens collected at Rjazanovka, the extreme South Primorje.

REFERENCES

- BANKS, N. (1905): A revision of the Nearctic Hemerobiidae. — Trans. Amer. Ent. Soc., **32**: 21—51.
- DOROKHOVA, G. I. (1979): Lacewings of the family Chrysopidae (Neuroptera) of the USSR. — Ent. Obozr., **58**: 105—111. (in Russian.)
- DOROKHOVA, G. I. (1980): Chrysopa perla (Linnaeus, 1758) (Neuroptera, Chrysopidae). — In: The insect areas of the USSR European part, Nauka Publ., Leningrad, p. 21 (map 38) (in Russian).

- DOROKHOVA, G. I. (1981): *Chrysopa prasina* Burmeister, 1839 (Neuroptera, Chrysopidae). — In: The insect areas of the USSR, European part, Nauka Publ., Leningrad, p. 5. (map 74) (in Russian).
- DOROKHOVA, G. I. (1987): Neuropteroidea. — In: Identification book of insects of the USSR, European part, Nauka Publ., Leningrad, vol. 4, pt. 6, p. 7—96 (in Russian).
- ESBEN-PETERSEN, P. (1921): Notes concerning some Neuroptera in the Helsingfors Museum together with a description of *Hemerobius poppii*. — Nor. Ent., **1**: 38—43.
- GERSTAECKER, A. (1885): Vier Decaden von Neuropteren aus der Familie Megaloptera Burm. — Mitt. naturwiss. Ver. Neuvorpomm. Rügen, **16**: 1—49.
- HAGEN, H. A. (1858): Russlands Neuropteren. — Stett. Ent. Z., **19**: 110—134.
- HAGEN, H. A. (1866): *Hemerobidarum synopsis synonymica*. — Stett. Ent. Z., **27**: 369—462.
- HÖLZEL, H. (1970a): Myrmeleontidae aus den westlichen Teilen der Mongolie (Neuroptera-Planipennia). — Mitt. Zool. Mus. Berlin, **46**: 247—264.
- HÖLZEL, H. (1970b): Ergebnisse der zoologischen Forschungen von Dr. Z. Kaszab in der Mongolei. 222. Beitrag zur Kenntnis der Myrmeleontiden der Mongolei (Neuroptera-Planipennia). — Acta Zool. Hung., **16**: 115—136.
- KLIMASZEWSKI, J. & KEVAN, D. K. Mc. E. (1987): The brown lacewing flies of Canada and Alaska (Neuroptera: Hemerobiidae). Part. 2. The genus *Wesmaelius* Krüger. — Neur. Intern., **4**: 153—204.
- KOZHANTCHIKOV, I. V. (1953): Neuroptera. — In: Animal Kingdom of the USSR, Nauka Publ., Moscow & Leningrad, vol. 4, p. 424—433. (in Russian.)
- KUWAYAMA, S. (1924): A contribution to the knowledge of the Neuropterous insects of South Saghalien. — Trans. Sapporo—Nat. Hist. Soc., **9** (1): 93—140. (in Japanese.)
- KUWAYAMA, S. (1936): Materials for the study of the Neuropteroid fauna of the Kurile Islands. I. — Ins. Mats., **10**: 107—110.
- KUWAYAMA, S. (1954): Neuroptera-Planipennia from the Daisetsuzan National Park, Hokkaido, Japan. — Ins. Mats., **18**: 94—102.
- KUWAYAMA, S. (1956): Further studies on the Neuroptera-Planipennia of the Kurile Islands. — Ins. Mats., **20**: 77—82.
- KUWAYAMA, S. (1962): Revisional synopsis of the Neuroptera in Japan. — Pac. Ins., **4**: 325—412.
- KUWAYAMA, S. (1967): Insect fauna of the Southern Kurile Islands, Sapporo, 225 pp. (in Japanese.)
- LERAUT, P. (1980): Liste des Planipennes de France. — Bull. Soc. Ent. Fr., **85**: 237—253.
- MAKARKIN, V. N. (1984): The ant lions (Neuroptera, Myrmeleontidae) of the Far East. — In: Systematics of insects of the Far East, Far East Sci. Center Publ., Vladivostok, p. 38—40. (in Russian.)
- MAKARKIN, V. N. (1985a): A review of Hemerobiidae (Neuroptera) of the fauna of the USSR. I. The genera *Hemerobius* L., *Micromus* Ramb. and *Paramicromus* Nakah. — Ent. Obozr., **64**: 158—170. (in Russian.)
- MAKARKIN, V. N. (1985b): A contribution of the fauna of the Neuroptera from the Far East. — Zool. Zhurn., Moscow, **64**: 620—622. (in Russian.)
- MAKARKIN, V. N. (1985c): A review of the family Osmylididae (Neuroptera) of the USSR fauna. — In: Taxonomy and ecology of Arthropoda of the Far East, Far Eastern Sci. Center Publ., Vladivostok, p. 33—47. (in Russian.)
- MAKARKIN, V. N. (1985d): New or little known species of the green lacewings (Neuroptera, Chrysopidae) from the Far East. — In: Taxonomy and ecology of Arthropoda of the Far East, Vladivostok, p. 48—52. (in Russian.)
- MAKARKIN, V. N. (1985e): Influence of the conditions of a large city upon species composition in Neuroptera. — Ekologia, Sverdlovsk, N 4: 90—92. (in Russian.)
- MAKARKIN, V. N. (1985f): A faunal review of the green lacewings (Neuroptera, Chrysopidae) of the Far East. — In: Fauna and ecology of insects of the Primorje and Kamchatka (pests and entomophages), Far Eastern Sci. Center Publ., Vladivostok, p. 55—64. (in Russian.)
- MAKARKIN, V. N. (1986): A review of Hemerobiidae (Neuroptera) of the fauna of the USSR. 2. The genera *Wesmaelius* Krüger, *Symppherobius* Banks, *Psectra* Hagen, *Megalomus* Rambur, *Neuronema* McLach., and *Drepanepteryx* Leach. — Ent. Obozr., **65**: 604—617. (in Russian.)
- MATSUMURA, S. (1911): Erster Beitrag zur Insecten-Fauna von Saghalien. — J. Coll. Agr. Tohoku Imp. Univ., **4**: 1—144.
- MEINANDER, M. (1962): The Neuroptera and Mecoptera of Eastern Fennoscandia. — Fauna fenn., **13**: 1—96.

- MEINANDER, M. (1981): A review of the genus *Coniopteryx* (Neuroptera, Coniopterygidae). — Ann. Ent. Fenn., **47**: 97—110.
- NAKAHARA, W. (1956): New or little known Hemerobiidae from Japan and adjacent territories (Neuroptera). — Kontyû, **24**: 182—191.
- NAKAHARA, W. (1971): Some genera and species of the Hemerobiidae (Neuroptera). — Kontyû, **39**: 7—14.
- NAVÁS, L. (1912): Quelques Neuroptères de la Sibérie méridionale-orientale. — Rev. Ent. d'Rus., **12**: 414—422.
- NAVÁS, L. (1925): Entomologische Ergebnisse der schwedischen Kamtchatka-Expedition 1920—1922. 6. Neuroptera et Mecoptera. — Ark. Zool., **18** B: 1—4.
- NAVÁS, L. (1929): Insecta orientalia. VI—VII series. — Mem. Pont. Accad. Sci, Nuovi Lincei, sér 2, **12**: 33—56.
- OKAMOTO, H. (1926): Some Myrmeleonidae and Ascalaphidae from Corea. — Ins. Mats., **1**: 18—23.
- PLESHANOV, A. S. (1974): A review of the lacewings (Neuroptera) of Siberia and Far East. — In: Fauna of insects of the Eastern Siberia and Far East, Irkutsk State Univ. Publ., p. 180—193 (in Russian.)
- SHUVAKHINA, E. YA. (1980a): *Crysopa formosa* Brauer, 1850. — In: The insects areas of the USSR European part, Nauka Publ., Leningrad, p. 22. (map 39) (in Russian).
- SHUVAKHINA, E. YA. (1980b): *Chrysopa septempunctata* Wesmael, 1841. — In: The insect areas of the USSR European part, Nauka Publ., Leningrad, p. 23 (map 40). (in Russian.)
- TJEDER, B. (1936): Schwedisch-chinesische wissenschaftliche Expedition nach den nord-westlichen Provinzen Chinas. 62. Neuroptera. — Ark. Zool., **28** A: 1—36.
- TSUKAGUCHI, T. (1985): A check-list of published species of Japanese Chrysopidae (Neuroptera). — Kontyû, **53**: 503—506.
- ZAKHARENKO, A. V. (1987): Neuroptera of the fauna of the USSR. I. Family Mantispidae. — Ent. Obozr., **66**: 621—626 (in Russian).
- ZAKHARENKO, A. V. (1987b): A new species of the genus *Coniopteryx* (Neuroptera, Coniopterygidae) from the Far East. — Vestn. Zool., Kiev, N **5**: 76 (in Russian).
- ZAKHARENKO, A. V. (1988): Neuroptera of the fauna of the USSR. II. Fam. Dilaridae, Berothidae and Sisyridae. — Ent. Obozr., **67**: 763—768 (in Russian).
- ZAKHARENKO, A. V. & SEDYKH, K. F. (1981): Neuroptera of the Komi ASSR. — Ent. Obozr., 598—600 (in Russian))

Bibliography of the Neuropterida

Bibliography of the Neuropterida Reference number (r#):
7117

Reference Citation:

Makarkin, V. N. 1990 [1990.??.??]. A check-list of the Neuroptera-Planipennia of the USSR far east, with some taxonomic remarks. *Acta Zoologica Hungarica* 36:37-45.

Copyrights:

Any/all applicable copyrights reside with, and are reserved by, the publisher(s), the author(s) and/or other entities as allowed by law. No copyrights belong to the Bibliography of the Neuropterida. Work made available through the Bibliography of the Neuropterida with permission(s) obtained, or with copyrights believed to be expired.

Notes:

File:

File produced for the Bibliography of the Neuropterida (BotN) component of the Lacewing Digital Library (LDL) Project, 2009.